Fault-free Hamiltonian cycles in balanced hypercubes with conditional edge faults

Pingshan Li Min Xu*

Sch. Math. Sci. & Lab. Math. Com. Sys., Beijing Normal University, Beijing, 100875, China

arXiv:1701.03216v1 [math.CO] 12 Jan 2017

Abstract

The balanced hypercube, BH_n, is a variant of hypercube Q_n. Zhou et al. [Inform. Sci. 300 (2015) 20-27] proposed an interesting problem that whether there is a fault-free Hamiltonian cycle in BH_n with each vertex incident to at least two fault-free edges. In this paper, we consider this problem and show that each fault-free edge lies on a fault-free Hamiltonian cycle in BH_n after no more than $4n-5$ faulty edges occur if each vertex is incident with at least two fault-free edges for all $n \geq 2$. Our result is optimal with respect to the maximum number of tolerated edge faults.

Key words: Balanced hypercubes; Hypercubes; Hamiltonian cycles; Fault-tolerance.

1 Introduction

In the field of parallel and distributed systems, interconnection networks are an important research area. Typically, the topology of a network can be represented as a graph in which the vertices represent processors and the edges represent communication links.

For graph definitions and notations, we follow [1]. A graph G consists of a vertex set $V(G)$ and an edge set $E(G)$, where an edge is an unordered pair of distinct vertices of G. A graph G is called bipartite if its vertex set can be partitioned into two parts V_1, V_2 such that every edge has one endpoint in V_1 and one in V_2. A vertex v is a neighbor of u if (u, v) is an edge of G, and $N_G(u)$ denotes all the neighbors of u in G. A path P of length ℓ from x to y is a finite sequence of distinct vertices $\langle v_0, v_1, \cdots, v_\ell \rangle$ such that $x = v_0, y = v_\ell$, and $(v_i, v_{i+1}) \in E$ for $0 \leq i \leq \ell - 1$. We also denote the path P as $\langle v_0, v_1, \cdots, v_i, Q, v_j, v_{j+1}, \cdots, v_\ell \rangle$, where Q is the path $\langle v_i, v_{i+1}, \cdots, v_j \rangle$. A cycle C of length $\ell+1$ is a closed path$\langle v_0, v_1, \cdots, v_\ell, v_0 \rangle$.

A path (resp., cycle) is called a Hamiltonian path (resp., cycle) if it contains all the vertices of G. A graph G is said to be Hamiltonian if there is a Hamiltonian cycle. A graph G is said to be Hamiltonian connected if there is a Hamiltonian path between any two vertices of G. A bipartite graph is Hamiltonian laceable if there is a Hamiltonian path between any two vertices in different bipartite sets.

The Hamiltonian property is one of the major requirements in designing network topologies since a topology structure containing Hamiltonian paths or cycles can efficiently simulate many algorithms that are designed on linear arrays or rings [16].

It is important to consider fault-tolerance in networks since faults may occur in real networks. Two fault models have been studied in many well-known networks, one is the random fault model, which means that faults may occur anywhere without any restriction, see, for example, [13, 14, 18, 24]. The other is the conditional fault model, which assumes that the fault distribution is limited. For example, some studies on two or more non-faulty edges incident to each vertex can be found in [2, 7, 9, 10, 15, 19, 21].

The hypercube network has been proved to be one of the most popular interconnection networks because it possesses many excellent properties such as a recursive structure, regularity, and symmetry.

*Corresponding author.
E-mail address: xum@bnu.edu.cn (M. Xu).

The balanced hypercube, proposed by Huang and Wu [8], is a hypercube variant. Similar to hypercubes, balanced hypercubes are bipartite graphs [8] that are vertex-transitive [17] and edge-transitive [26]. Balanced hypercubes are superior to hypercubes in that they have a smaller diameter than hypercubes[17].

The balanced hypercube, BH_n, has been studied by many researchers in recent years. Xu et al. [20] proved that BH_n is edge-bipancyclic and Hamiltonian laceable. Yang [22] proved that BH_n is bipanconnected. Yang [23] also demonstrated that the super connectivity of BH_n is $4n-4$ and the super edge-connectivity of BH_n is $4n-2$ for $n \geq 2$. Lü et al. [12] proved that BH_n is hyper-Hamiltonian laceable. Cheng et al. [3] proved that BH_n is $(n-1)$-vertex-fault-tolerant edge-bipancyclic. Hao et al. [6] showed that there is a fault-free Hamiltonian path between any two adjacent vertices in BH_n with $2n-2$ faulty edges. Cheng et al. [5] proved that BH_n is $2n-3$ edge-fault-tolerant 6-edge-bipancyclic for all $n \geq 2$. Zhou et al. [25] proved that BH_n is $2n-2$ edge-fault-tolerant Hamiltonian laceable, and they proposed an interesting problem that whether there is a fault-free Hamiltonian cycle in BH_n with each vertex incident to at least two fault-free edges. In this paper, we consider this problem and show that each fault-free edge lies on a fault-free Hamiltonian cycle in BH_n after no more than $4n-5$ faulty edges occur if each vertex is incident with at least two fault-free edges for all $n \geq 2$. Our result is optimal with respect to the maximum number of tolerated edge faults.

The rest of this paper is organized as follows. In Section 2, we introduce two equivalent definitions of balanced hypercubes and discuss some of their properties. In section 3, we introduce some lemmas used in the proof of the main result and prove the main result. Finally, we conclude this paper and give an example to show that our result is optimal in Section 4 .

2 Balanced hypercubes

Wu and Huang [8] present two equivalent definitions of BH_n as follows:

Definition 2.1 *An n-dimensional balanced hypercube BH_n has 2^{2n} vertices, each labeled by an n-bit string $(a_0, a_1, \cdots, a_{n-1})$, where $a_i \in \{0,1,2,3\}$ for all $0 \leq i \leq n-1$. An arbitrary vertex $(a_0, a_1, \cdots, a_{i-1}, a_i, a_{i+1}, \cdots, a_{n-1})$ is adjacent to the following $2n$ vertices:*

(1) $((a_0 \pm 1) \mod 4, a_1, \cdots, a_{i-1}, a_i, a_{i+1}, \cdots, a_{n-1})$ *where* $1 \leq i \leq n-1$,
(2) $((a_0 \pm 1) \mod 4, a_1, \cdots, a_{i-1}, (a_i + (-1)^{a_0}) \mod 4, a_{i+1}, \cdots, a_{n-1})$ *where* $1 \leq i \leq n-1$.

In BH_n, the first coordinate a_0 of vertex $(a_0, a_1, \cdots, a_{n-1})$ is called the inner index, and the second coordinate $a_i (1 \leq i \leq n-1)$ is called the i-dimension index. From the definition, we have $N_{BH_n}((a_0, a_1, \cdots, a_{n-1})) = N_{BH_n}((a_0+2, a_1, \cdots, a_{n-1}))$. Figure 1 shows two balanced hypercubes of dimensional one and two.

Briefly, we assume that '$+,-$' for the coordinate of a vertex is an operation with mod 4 in the remainder of the paper. Let $X_{j,i} = \{(a_0, a_1, \cdots, a_{j-1}, a_j, a_{j+1}, \cdots, a_{n-1}) \mid a_k \in \{0,1,2,3\}, 0 \leq k \leq n-1, a_j = i\}$ for $1 \leq j \leq n-1$ and $i \in \{0,1,2,3\}$ and let $BH_{n-1}^{j,i} = BH_n[X_{j,i}]$. Then, BH_n can be divided into four copies: $BH_{n-1}^{j,0}, BH_{n-1}^{j,1}, BH_{n-1}^{j,2}$, and $BH_{n-1}^{j,3}$ where $BH_{n-1}^{j,i} \cong BH_{n-1}$ for $i = 0,1,2,3$ [3]. We use BH_{n-1}^{i} to denote $BH_{n-1}^{n-1,i}$ for $i = 0,1,2,3$.

Definition 2.2 *An n-dimensional balanced hypercube BH_n can be constructed recursively as follows:*

1. *BH_1 is a cycle of length four with vertex-set $\{0,1,2,3\}$.*
2. *BH_n is a construct from four copies of BH_{n-1} : $BH_{n-1}^0, BH_{n-1}^1, BH_{n-1}^2$, and BH_{n-1}^3. Each vertex $(a_0, a_1, \cdots, a_{n-2}, i)$ has two extra adjacent vertices:*
   (1) *In BH_{n-1}^{i+1} : $(a_0 \pm 1, a_1, \cdots, a_{n-2}, i+1)$ if a_0 is even.*
   (2) *In BH_{n-1}^{i-1} : $(a_0 \pm 1, a_1, \cdots, a_{n-2}, i-1)$ if a_0 is odd.*



Figure 1: Illustration of BH_1 and BH_2

Since BH_n is a bipartite graph, $V(BH_n)$ can be divided into two disjoint parts. Obviously, the vertex-set $V_1 = \{a = (a_0, a_1, \cdots, a_{n-1}) \mid a \in V(BH_n)$ with a_0 odd$\}$ and $V_2 = \{a = (a_0, a_1, \cdots, a_{n-1}) \mid a \in V(BH_n)$ with a_0 even$\}$ form the desired partition. We use black nodes to denote the vertices in V_1 and white nodes to denote the vertices in V_2.

Let (u, v) be an edge of BH_n. If u and v differ only with regard to the inner index, then (u, v) is said to be a 0-dimension edge. If u and v differ not only in terms of the inner index but also with regard to the i-dimension index, then (u, v) is called the i-dimension edge. We use $\partial D_d (0 \leq d \leq n-1)$ to denote the set of all d-dimension edges.

There are some known properties about BH_n.

Lemma 2.3 ([17, 26]) *The balanced hypercube BH_n is vertex-transitive and edge-transitive.*

A bipartite graph G is k-fault-tolerant Hamiltonian laceable if $G - F$ remains Hamiltonian laceable for $F \subseteq V(G) \cup E(G)$ with $|F| \leq k$. A bipartite graph G is k-edge-fault-tolerant Hamiltonian laceable if $G - F$ remains Hamiltonian laceable for $F \subseteq E(G)$ with $|F| \leq k$. Zhou et al. obtained the following result.

Lemma 2.4 ([25]) *The balanced hypercube BH_n is $(2n-2)$-edge-fault-tolerant Hamiltonian laceable for $n \geq 2$.*

Lemma 2.5 ([11]) *Let $n \geq 2$. Then, $BH_n - \partial D_0$ has four components, and each component is isomorphic to BH_{n-1}.*

The above lemma shows that one can divide BH_n into four BH_ns by deleting ∂D_d for any $d \in \{0, 1, \cdots, n-1\}$. The four components of $BH_n - \partial D_j$ are $BH_{n-1}^{j,0}$, $BH_{n-1}^{j,1}$, $BH_{n-1}^{j,2}$, and $BH_{n-1}^{j,3}$ for $1 \leq j \leq n-1$. For convenience, we use $BH_{n-1}^{0,0}, BH_{n-1}^{0,1}, BH_{n-1}^{0,2}$, and $BH_{n-1}^{0,3}$ to denote the components of $BH_n - \partial D_0$ throughout this paper.

Lemma 2.6 ([4]) *Let U, V be two distinct bipartitions of BH_n, u^1, u^2 be two different vertices in U, and v^1, v^2 be two different vertices in V. Then, there are two disjoint paths P, Q such that* (1) *P joins u^1 and v^1, and Q joins u^2 and v^2;* (2) *$V(P \cup Q) = V(BH_n)$.*

A graph G is hyper-Hamiltonian laceable if it is Hamiltonian laceable and, for an arbitrary vertex v in V_i where $i \in \{0, 1\}$, there is a Hamiltonian path in $G - v$ joining any two different vertices in V_{1-i}. Lü et al. obtained the following result.

Lemma 2.7 ([12]) *The balanced hypercube BH_n is hyper-Hamiltonian laceable for $n \geq 1$.*

3 Main result

In this section, we will give the proof of the main result. First, we introduce some lemmas that will be used in the proof of the main result.

Definition 3.1 *Suppose that G is a graph and $F \subseteq E(G)$. A vertex u is called i-rescuable in G if $|N_{G-F}(u)| = i$.*

Lemma 3.2 *Let $n \geq 3, F \subseteq E(BH_n)$ with $|F| = 4n-5$ and $\delta(BH_n - F) \geq 2$. Then, at least one statement holds in the following* (1) *and* (2).

(1) *There is an integer m in $\{0, 1, \cdots, n-1\}$ such that $|F \cap \partial D_m| \geq 3$ and $\delta(BH_n - \partial D_m) \geq 2$.*

(2) *There are two integers m, m' in $\{0, 1, \cdots, n-1\}$ such that $|F \cap \partial D_m| \geq 2$ and $|F \cap \partial D_{m'}| \geq 2$. Furthermore, there is no isolated vertex and no more than one 1-rescuable vertex in $BH_n - \partial D_m$ (resp., $BH_n - \partial D_{m'}$).*

Proof: Let F be a faulty set in $E(BH_n)$ with $|F| = 4n-5$ and $\delta(BH_n - F) \geq 2$. Suppose that u is k-rescuable, and v is t-rescuable in BH_n where $k \leq t \leq \min\{m \mid x \text{ is } m\text{-rescuable in } BH_n, \text{ where } x \in V(BH_n) \setminus \{u, v\}\}$.

If $k \geq 4$, there is an integer $m \in \{0, 1, \cdots, n-1\}$ such that $|F \cap \partial D_m| \geq 3$ because $|F| = 4n-5 > 2n$. Each vertex is at least 2-rescuable in $BH_n - \partial D_m$ for $k \geq 4$. Thus, statement (1) holds. In the following, we assume that $2 \leq k \leq 3$ and suppose that the dimensions of fault-free edges incident with u are $i_1, i_2, \cdots, i_k$ and the dimensions of fault-free edges incident with v are $j_1, j_2, \cdots, j_t$ where $\{i_1, i_2, \cdots, i_k\}$ and $\{j_1, j_2, \cdots, j_t\}$ may be multisets.

Case 1: $k = 2$.

Subcase 1.1: $t = 2$.

In this subcase, $|F \cap \{(u, x) \mid x \in N_{BH_n}(u)\}| = |F \cap \{(v, x) \mid x \in N_{BH_n}(v)\}| = 2n-2$. Since $2(2n-2) - (4n-5) = 1$, we obtain $(u, v) \in F$, and there is no faulty edge in $BH_n - \{u, v\}$. Note that there is no triangle in BH_n. For an arbitrary vertex x in $V(BH_n) \setminus \{u, v\}$, x is incident with no more than one faulty edge, which means that x is at least 3-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$ as $2(n-1) - 1 \geq 3$. Note that for $4n-5 > 2n$, there is an $m \in \{0, 1, \cdots, n-1\}$ such that $|F \cap \partial D_m| \geq 3$.

If $m \notin \{i_1, i_2, j_1, j_2\}$, then u and v are 2-rescuable in $BH_n - \partial D_m$. Therefore, statement (1) holds.

If $m \in \{i_1, i_2, j_1, j_2\}$, without loss of generality, let $i_1 = m$. Note that each vertex in BH_n is incident with two i-dimension edges for $i \in \{0, 1, \cdots, n-1\}$. The integer m appears no more than once in the multiset $\{i_1, i_2, j_1, j_2\}$. Thus, u is 1-rescuable, and v is 2-rescuable in $BH_n - \partial D_m$. Note that for $n \geq 3$, according to the pigeonhole principle, there is an integer $m' \in \{0, 1, \cdots, n-1\} \setminus \{m\}$ such that m' appears no more than once in the multiset $\{i_2, j_1, j_2\}$. Thus, $|F \cap \partial D_{m'}| \geq 2$. Without loss of generality, we can assume that u is 1-rescuable and v is 2-rescuable in $BH_n - \partial D_{m'}$. Therefore, statement (2) holds.

Subcase 1.2: $t = 3$.

First, we prove that each vertex in $V(BH_n) \setminus \{u, v\}$ is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$.

If $(u, v) \notin F$, there are $4n-5$ faulty edges incident with u or v. This means there is no faulty edge in $BH_n - \{u, v\}$. Thus, each vertex in $V(BH_n) \setminus \{u, v\}$ is incident with no more than two faulty edges. That means it is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$ since $2(n-1) - 2 \geq 2$.

If $(u, v) \in F$, there are $4n-6$ faulty edges incident with u or v. This means that there is no more than one faulty edge in $BH_n - \{u, v\}$. Since there is no triangle in BH_n, each vertex in $V(BH_n) \setminus \{u, v\}$ is incident with no more than two faulty edges. That implies that it is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$, as $2(n-1) - 2 \geq 2$.

Next, we complete the proof of subcase 1.2.

If there is an $m \in \{0, 1, \cdots, n-1\}$ such that $m \notin \{i_1, i_2, j_1, j_2, j_3\}$, then $|F \cap \partial D_m| \geq 3$ and u and v are at least 2-rescuable in $BH_n - \partial D_m$. Therefore, statement (1) holds.

Now, we assume that $\{0, 1, \cdots, n-1\} \subseteq \{i_1, i_2, j_1, j_2, j_3\}$. If $n \geq 4$ or $n = 3$ and $i_1 = i_2$, there are two distinct integers m, m' in $\{0, 1, \cdots, n-1\} \setminus \{i_1, i_2\}$. Note that u is incident with two faulty

m (resp., m')-dimension edges and u is 2-rescuable, v is at least 1-rescuable in $BH_n - \partial D_m$ (resp., $BH_n - \partial D_{m'}$). Therefore, statement (2) holds.

If $n = 3$ and $i_1 \neq i_2$, without loss of generality, let $i_1 = 0, i_2 = 1$. Then $2 \in \{j_1, j_2, j_3\}$. Thus, u is 2-rescuable, and v is at least 1-rescuable in $BH_n - \partial D_2$. Let F' be the set of faulty edges incident with u or v. Then $|F'| \geq 4n - 6 = 6$ and $|F \cap \partial D_2| \geq |F' \cap \partial D_2| \geq 2$. Moreover, $2 \leq |F' \cap \partial D_2| \leq 3$. Further, we obtain $|F' \cap \partial D_0| \geq 2$ or $|F' \cap \partial D_1| \geq 2$, as $|F'| \geq 4n - 6 = 6$. Without loss of generality, we can assume that $|F' \cap \partial D_0| \geq 2$. Since u is incident with exactly one faulty 0-dimension edge, v is incident with at least one faulty 0-dimension edge. Thus, 0 appears no more than once in the multiset $\{j_1, j_2, j_3\}$. Note that u is 1-rescuable, v is at least 2-rescuable in $BH_n - \partial D_0$ and $|F \cap \partial D_0| \geq |F' \cap \partial D_0| \geq 2$. Let $m = 0, m' = 2$. Therefore, statement (2) holds.

Subcase 1.3: $t \geq 4$.

Since $t \geq 4$, each vertex in $BH_n - \{u\}$ is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$. Suppose that there are two distinct integers $m, m' \in \{0, 1, \cdots, n-1\} \setminus \{i_1, i_2\}$. Note that u is incident with two faulty m (resp., m')-dimension faulty edges, and u is 2-rescuable in $BH_n - \partial D_m$ (resp., $BH_n - \partial D_{m'}$). Then, statement (2) holds.

If there aren't two distinct integers $m, m' \in \{0, 1, \cdots, n-1\} \setminus \{i_1, i_2\}$, then $n = 3, i_1 \neq i_2$ and $|F| = 4n - 5 = 7$. Without loss of generality, we can assume that $i_1 = 0, i_2 = 1$. Then, $|F \cap \partial D_2| \geq 2$. If $|F \cap \partial D_2| \geq 3$, then u is 2-rescuable in $BH_n - \partial D_2$. Therefore, statement (1) holds. If $|F \cap \partial D_2| = 2$, then u is 2-rescuable in $BH_n - \partial D_2$. According to the pigeonhole principle, $|F \cap \partial D_0| \geq 3$ or $|F \cap \partial D_1| \geq 3$. Without loss of generality, we can assume that $|F \cap \partial D_0| \geq 3$. Thus, u is 1-rescuable in $BH_n - \partial D_0$. Let $m = 0, m' = 2$. Therefore, statement (2) holds.

Case 2: $k = 3$.

Subcase 2.1: $t = 3$.

(a) There is a 3-rescuable vertex w in BH_n where $w \in V(BH_n) \setminus \{u, v\}$.

Suppose that the dimensions of fault-free edges incident with w are s_1, s_2, s_3. Note that $3(2n-3) - (4n-5) = 2n - 4 \geq 2$ and there is no triangle in BH_n. We obtain $n = 3$ and $|E(BH_n[\{u, v, w\}])| = |F \cap E(BH_n[\{u, v, w\}])| = 2$. Without loss of generality, let $(u, v), (v, w) \in F$. Since $(4 \times 3 - 5) - (6 \times 3 - 11) = 0$, $F \subseteq \{e \mid e \text{ is incident to a vertex in } \{u, v, w\}\}$. Hence, each vertex in $V(BH_n) \setminus \{u, v, w\}$ is incident with no more than two faulty edges. This means that each vertex in $V(BH_n) \setminus \{u, v, w\}$ is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$.

Note that $4n - 5 = 7 > 2n$. There is an integer $m \in \{0, 1, \cdots, n-1\}$ such that $|F \cap \partial D_m| \geq 3$. Without loss of generality, we can assume that $m = 0$. Since each vertex is incident with exactly two 0-dimension edges, 0 appears no more than three times in the multiset $\{i_1, i_2, i_3, j_1, j_2, j_3, s_1, s_2, s_3\}$.

If 0 does not appear twice in the multiset $\{i_1, i_2, i_3\}$, $\{j_1, j_2, j_3\}$ and $\{s_1, s_2, s_3\}$, then u, v, w are at least 2-rescuable in $BH_n - \partial D_0$. Thus, statement (1) holds.

Otherwise, without loss of generality, assume that $i_1 = i_2 = 0, i_3 = 1$. Then, u is 1-rescuable, and v and w are at least 2-rescuable in $BH_n - \partial D_0$. Moreover, u is 2-rescuable in both $BH_n - \partial D_1$ and $BH_n - \partial D_2$.

If 2 appears four times in the multiset $\{j_1, j_2, j_3, s_1, s_2, s_3\}$, then $|F \cap \partial D_1| \geq 2$. Note that v, w are at least 2-rescuable in $BH_n - \partial D_1$. Then, let $m = 0, m' = 1$. Statement (2) holds.

If 2 appears no more than three times in the multiset $\{j_1, j_2, j_3, s_1, s_2, s_3\}$, then one of the vertices in $\{v, w\}$ is at least 2-rescuable, and another is at least 1-rescuable in $BH_n - \partial D_2$. Let $m = 0, m' = 2$. Statement (2) holds.

(b) Suppose that each vertex in $V(BH_n) \setminus \{u, v\}$ is at least 4-rescuable in BH_n.

Note that each vertex in $V(BH_n) \setminus \{u, v\}$ is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$. Let F' be the set of faulty edges incident with u or v. Then, $|F'| \geq 4n - 7$.

If there is an integer $m \in \{0, 1, \cdots, n-1\}$ such that $|F' \cap \partial D_m| \geq 3$, then m appears no more than once in the multiset $\{i_1, i_2, i_3, j_1, j_2, j_3\}$. Thus, u (resp., v) is at least 2-rescuable in $BH_n - \partial D_m$. Statement (1) holds.

Otherwise, $n = 3$ and $|F' \cap \partial D_i| \leq 2$ for all $i = 0, 1, 2$. Then, there are two distinct integers $m, m' \in \{0, 1, 2\}$ such that (2) holds. In fact, there are two integers $m, m' \in \{0, 1, 2\}$ such that $|F' \cap \partial D_m| = 2$ and $|F' \cap \partial D_{m'}| = 2$, as $|F'| \geq 4n - 7 = 5$. Thus, m (resp., m') appears no more

than twice in the multiset $\{i_1, i_2, i_3, j_1, j_2, j_3\}$. Hence, one vertex in $\{u, v\}$ is at least 2-rescuable and another is at least 1-rescuable in $BH_n - \partial D_m$ (resp., $BH_n - \partial D_{m'}$). Statement (2) holds.

Subcase 2.2: $t \geq 4$.

Note that for $t \geq 4$, each vertex in $BH_n - \{u\}$ is at least 2-rescuable in $BH_n - \partial D_i$ for all $i \in \{0, 1, \cdots, n-1\}$. Since $4n - 5 > 2n$, there is an integer m such that $|F \cap \partial D_m| \geq 3$. If m appears no more than once in the multiset $\{i_1, i_2, i_3\}$, then u is at least 2-rescuable in $BH_n - \partial D_m$. Statement (1) holds. If m appears twice in the multiset $\{i_1, i_2, i_3\}$, without loss of generality, let $i_1 = i_2 = m$. Then, u is 1-rescuable in $BH_n - \partial D_m$ and let m' be an integer in $\{0, 1, \cdots, n-1\} \setminus \{m, i_3\}$. Then u is incident with two faulty m'-dimension edges and u is 3-rescuable in $BH_n - \partial D_{m'}$. Hence, statement (2) holds.

Owing to the above discussion, the lemma holds. □

Lemma 3.3 *Let $n \geq 3, F \subseteq E(BH_n)$ with $|F| \leq 4n - 5$. Then, for any vertex u in $BH_{n-1}^{j,i}$, there is a fault-free j-dimension edge (v, w) such that $(u, v) \in E(BH_{n-1}^{j,i})$. Moreover, if u is 1-rescuable in $BH_n - \partial D_j$, then (u, v) can be a faulty edge.*

Proof: Each vertex in $V(BH_n)$ is incident with two i-dimension edges for all $i \in \{0, 1, \cdots, n-1\}$. Thus, for any vertex $u \in BH_{n-1}^{j,i}$, its degree is $2n - 2$ in $BH_{n-1}^{j,i}$. There is at least one fault-free j-dimension edge (v, w) such that $(u, v) \in E(BH_{n-1}^{j,i})$ for $2(2n-2) - (4n-5) \geq 1$. If u is incident with only one fault-free edge in $BH_{n-1}^{j,i}$, then u is incident with $2n - 3$ faulty edges in $BH_{n-1}^{j,i}$. Thus, $|F \cap \partial D_j| \leq 2n - 2$. Since $2(2n-3) - (2n-2) = 2n - 4 \geq 2$, there is at least one fault-free edge (w, v) such that $(u, v) \in F$.

□

Lemma 3.4 *Let $F \subseteq E(BH_2)$ with $|F| \leq 3$ and $\delta(BH_2 - F) \geq 2$. Then, each edge in $BH_2 - F$ lies on a fault-free Hamiltonian cycle.*

Proof: The proof is rather long, so we provide it in Appendix A. □

Theorem 3.5 *Let $F \subseteq E(BH_n)$ with $|F| \leq 4n-5$ and $\delta(BH_n - F) \geq 2$. Then, each edge in $BH_n - F$ lies on a fault-free Hamiltonian cycle.*

Proof: We prove this theorem by induction on n. By Lemma 3.4, the theorem holds for $n = 2$. Assume that this is true for $2 \leq k \leq n-1$. By Lemma 3.2, one may partite BH_n along dimension j, $0 \leq j \leq n-1$, into four $BH_{n-1}s$, denoted by $BH_{n-1}^{j,0}, BH_{n-1}^{j,1}, BH_{n-1}^{j,2}$ and $BH_{n-1}^{j,3}$, such that $|F \cap \partial D_j| \geq 2$ and there is no isolated vertex and no more than one 1-rescuable vertex in $BH_n - \partial D_j$. Let $F^i = F \cap BH_{n-1}^{j,i}$ and $e = (u, v)$ be an arbitrary edge in $BH_n - F$. We must show that there is a Hamiltonian cycle in $BH_n - F$ that contains e.

Case 1: $e \in E(BH_n) - \partial D_j$.

Without loss of generality, we can assume that $e \in BH_{n-1}^{j,0}$.

Subcase 1.1: $|F^i| \leq 4n - 9$ for all $i = 0, 1, 2, 3$.

Subcase 1.1.1: $\delta(BH_n - F - \partial D_j) = 1$.

By Lemma 3.2, there is exactly one 1-rescuable vertex in $BH_n - \partial D_j$, say, w. Without loss of generality, we can assume that w is a white vertex.

Subcase 1.1.1.1: $w \in BH_{n-1}^{j,0}$.

Since $\delta(BH_n - F) \geq 2$, w is incident with at least one fault-free j dimension edge, say, (w, b^1). By Lemma 3.3, there is a fault-free j-dimension edge (b^0, a^3) such that $(w, b^0) \in F$. By induction, there is a Hamiltonian cycle C_0 in $BH_{n-1}^{j,0} - F + (w, b^0)$ that contains (u, v). Since w is incident with exactly two edges in $BH_{n-1}^{j,0} - F + (w, b^0)$, $(w, b^0) \in E(C_0)$. We represent C_0 as $\langle u, H_0, w, b^0, H_0', u \rangle$. Note that w is incident with $2n - 3$ faulty edges in $BH_{n-1}^{j,0}$, so we have $|F^0| \geq 2n - 3$. Hence $|F^1| + |F^2| + |F^3| = |F| - |F^0| - |F \cap \partial D_j| \leq 4n - 5 - (2n - 3) - 2 = 2n - 4$. Let $(a^1, b^2), (a^2, b^3)$ be fault-free j-dimension edges. By Lemma 2.4, there is a fault-free Hamiltonian path H_i in $BH_{n-1}^{j,i}$ that



Figure 2: Illustration for subcase 1.1.1.1 of theorem 3.5

joins a^i and b^i for $i = 1, 2, 3$. Then, the cycle $C = \langle u, H_0, w, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H'_0, u\rangle$ (see figure 2) is the desired cycle.

Subcase 1.1.1.2: $w \in BH^{j,1}_{n-1}$ (or $BH^{j,3}_{n-1}$).

Since $4n-5-(2n-3)-2 = 2n-4$, $|F^i| \leq 2n-4$ for all $i = 0, 2, 3$.

Let $X = \{x^0 \mid (x^0, b^1) \in \partial D_j \setminus F$, where $x^0 \in BH^{j,0}_{n-1}$ and $(w, b^1) \in F\}$. Since $2(2n-3) - [(4n-5)-(2n-3)] = 2n-4 \geq 2$, we can obtain $|X| \geq 1$. Moreover, if $|X| = 1$, there are $2n-2$ faulty edges between $BH^{j,0}_{n-1}$ and $BH^{j,1}_{n-1}$.

Subcase 1.1.1.2.1: $X = \{u\}$.

By induction, there is a fault-free Hamiltonian cycle C_0 in $BH^{j,0}_{n-1}$ that contains (u, v) and a Hamiltonian cycle C_1 in $BH^{j,1}_{n-1} - F + (w, b^1)$ that contains (w, b^1). Suppose that $N_{C_0}(u) = \{v, b^0\}$. Let $H_0 = C_0 - (u, b^0)$, $H_1 = C_1 - (w, b^1)$. Note that $|X| = 1$, so there are $2n-2$ faulty j-dimension edges between $BH^{j,0}_{n-1}$ and $BH^{j,1}_{n-1}$. Thus, $F \cap E(BH^{j,0}_{n-1}, BH^{j,3}_{n-1}) = \emptyset$. Let $(b^0, a^3), (a^2, b^3)$ be fault-free j-dimension edges. Note that $|F^i| \leq 2n-4$ for $i = 2, 3$. By Lemma 2.4, there is a Hamiltonian path H_i of $BH^{j,i}_{n-1}$ that joins a^i and b^i for $i = 2, 3$. Hence, the cycle $C = \langle u, b^1, H_1, w, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 3) is the desired cycle.

Subcase 1.1.1.2.2: $X \neq \{u\}$.

(a) $|F \cap \partial D_j| \leq 3$.

Let $a^0 \in X \setminus \{u\}$ and (w, b^2) be a fault-free j-dimension edge. By induction, there is a fault-free Hamiltonian cycle C_0 in $BH^{j,0}_{n-1}$ that contains (u, v). Suppose that $N_{C_0}(a^0) = \{b^0, d^0\}$. Note $|F \cap \partial D_j| \leq 3$; we can see that b^0 or d^0 is incident with one fault-free j-dimension edge. Without loss of generality, we can assume that b^0 is incident with one fault-free j-dimension edge, say, (b^0, a^3). We can represent C_0 as $\langle u, H_0, a^0, b^0, H'_0, u\rangle$. Let (a^2, b^3) be a fault-free j-dimension edge. Note that $|F^i| \leq (4n-5)-(2n-3)-2 = 2n-4$ for $i = 2, 3$. By Lemma 2.4, there is a fault-free Hamiltonian path H_i in $BH^{j,i}_{n-1}$ that joins a^i and b^i for $i = 2, 3$. By induction, there is a fault-free Hamiltonian cycle C_1 in $BH^{j,1}_{n-1}$ that contains (w, b^1) as $|F^1| \leq 4n-9$ and $\delta(BH^{j,1}_{n-1} - F + (w, b^1)) \geq 2$. Let $H_1 = C_1 - (w, b^1)$. Then, the cycle $C = \langle u, H_0, a^0, b^1, H_1, w, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H'_0, u\rangle$ (see figure 3) is the desired cycle.

(b) $|F \cap \partial D_j| \geq 4$.

Since $[2(n-1)-2]+4+(2n-3) = 4n-3 > 4n-5$, each vertex is at least 3-rescuable in $BH^{j,0}_{n-1}$. Let (u, α) be a fault-free edge in $BH^{j,0}_{n-1}$ where $\alpha \neq v$. Let $(F^0)' = F^0 \cup T$ where $T = \{(u, x) \mid x \in N_{BH^{j,0}(u)_{n-1}} - \{v, \alpha\}\}$. Then, $|(F^0)'| \leq 2n-4+[4n-5-(2n-3)-4] = 4n-10$, and $\delta(BH^{j,0}_{n-1} - F'_0) \geq 2$.

Let $a^0 \in X \setminus \{u\}$. By Lemma 3.3, there is a fault-free j-dimension edge (b^0, a^3) such that $(a^0, b^0) \in E(BH^{j,0}_{n-1})$. By induction, there is a Hamiltonian cycle C_0 in $(BH^{j,0}_{n-1} - (F^0)') \cup (a^0, b^0)$ that contains



Figure 3: Illustration for Subcase 1.1.1.2 of theorem 3.5

(a^0, b^0). Since u is incident with exactly two fault-free edges in $BH^{j,0}_{n-1} - (F^0)'$, $e = (u, v) \in E(C_0)$. We represent C_0 as $\langle u, H_0, a^0, b^0, H'_0, u\rangle$. Also by induction, there is a Hamiltonian cycle in $BH^{j,1}_{n-1} - F + (w, b^1)$ that contains (w, b^1). Let $H_1 = C_1 - (w, b^1)$ and (a^2, b^3) be a fault-free j-dimension edge. Note that $|F^i| \leq 2n - 4$ for $i = 2, 3$. By Lemma 2.4, there is a Hamiltonian path H_i in $BH^{j,i}_{n-1}$ that joins a^i and b^i for $i = 2, 3$. Then, the cycle $C = \langle u, H_0, a^0, b^1, H_1, w, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H'_0, u\rangle$ (see figure 3) is the desired cycle.

Subcase 1.1.1.3: $w \in BH^{j,2}_{n-1}$.

Since $\delta(BH_n - F) \geq 2$, w is incident with a fault-free j-dimension edge (w, b^3). By Lemma 3.3, there is a fault-free j-dimension edge (a^1, b^2) such that $(w, b^2) \in F$. By induction, there is a fault-free Hamiltonian cycle C_0 in $BH^{j,0}_{n-1}$ that contains (u, v). We represent it as $\langle c^1, c^2, \cdots, c^{2^{2n-2}}, c^1\rangle$ with $c^1 = u, c^{2^{2n-2}} = v$. Let $M = \{(c^1, c^2), \cdots, (c^{2i-1}, c^{2i}), \cdots, (c^{2^{2n-2}-1}, c^{2^{2n-2}})\}$. Therefore, M is a set of 2^{2n-3} mutually disjoint edges. There is an edge (c^{2i-1}, c^{2i}) in M such that c^{2i-1} (resp., c^{2i}) is incident with a fault-free j-dimension edge, say, (c^{2i-1}, b^1) (resp., (c^{2i}, a^3)) because $2 \cdot 2^{2n-3} > 4n - 5$. We can represent C_0 as $\langle u, H_0, c^{2i-1}, c^{2i}, H'_0, u\rangle$.



Figure 4: Illustration for subcase 1.1.1.3 of theorem 3.5

Note that $|F^i| \leq 2n-4$ for $i = 1, 3$. By Lemma 2.4, there is a fault-free Hamiltonian path H_i in $BH^{j,i}_{n-1}$ that joins a^i and b^i for $i = 1, 3$. By induction, there is a Hamiltonian cycle C_2 in $BH^{j,2}_{n-1} - F + (w, b^2)$ that contains (w, b^2). Let $H_2 = C_2 - (w, b^2)$. Then, the cycle $C = \langle u, H_0, c^{2i-1}, b^1, H_1, a^1, b^2, H_2, w, b^3, H_3, a^3, c^{2i}, H'_0, u\rangle$ (see figure 4) is the desired cycle.

Case 1.1.2: $\delta(BH_n - F - \partial D_j) \geq 2$.

Since $3(2n-4) - (4n-7) = 2n-5 > 0$, there is an integer $i \in \{1, 2, 3\}$ such that $|F^i| \leq 2n-4$. Without loss of generality, we can assume that $|F^1| \leq 2n-4$. By induction, there is a fault-free Hamiltonian cycle C_0 in $BH^{j,0}_{n-1}$ that contains (u, v). Similar to the analysis of subcase 1.1.1.3, we can represent C_0 as $\langle u, H_0, c^{2i-1}, c^{2i}, H'_0, u\rangle$ where $(c^{2i-1}, b^1), (c^{2i}, a^3)$ are fault-free j-dimension edges. By Lemma 3.3, there is a fault-free j-dimension edge (b^3, a^2) such that $(b^3, a^3) \in BH^{j,3}_{n-1}$. In addition, by Lemma 3.3, there is a fault-free j-dimension edge (b^2, a^1) such that $(a^2, b^2) \in E(BH^{j,2}_{n-1})$. By induction, there is a Hamiltonian cycle C_i in $(BH^{j,i}_{n-1} - F) \cup (a^i, b^i)$ that contains (a^i, b^i) for $i = 2, 3$. Let $H_i = C_i - (a^i, b^i)$ for i=2, 3. By Lemma 2.4, there is a fault-free Hamiltonian path H_1 in $BH^{j,1}_{n-1}$ that joins a^1 and b^1. Hence, the cycle $\langle u, H_0, c^{2i-1}, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, c^{2i}, H'_0, u\rangle$ (see figure 5) is the desired cycle.



Figure 5: Illustration for subcase 1.1.2 of theorem 3.5

Subcase 1.2: There is an integer $i \in \{0, 1, 2, 3\}$ such that $|F^i| = 4n-8$.

If $\delta(BH^{j,i}_{n-1} - F^i) \geq 2$, there are at least two vertex-disjoint faulty edges for $(4n-8)-[2(n-1)-2] = 2n-4 \geq 1$. Note that $|F \cap \partial D_j| \leq 3$. Then, there is a faulty edge (u^i, v^i) such that u^i is incident with a fault-free j-dimension edge and v^i is incident with a fault-free j-dimension edge.

Suppose that $\delta(BH^{j,i}_{n-1} - F^i) = 1$. Note that there is no more than one 1-rescuable vertex in $BH^{j,i}_{n-1}$. Let u^i be the 1-rescuable vertex in $BH^{j,i}_{n-1}$. Since $\delta(BH_n - F) \geq 2$, u^i is incident with a fault-free j-dimension edge. By Lemma 3.3, there is a fault-free j-dimension edge (v^i, x) such that $(u^i, v^i) \in F^i$. Obviously, $\delta(BH^{j,i}_{n-1} - F^i + (u^i, v^i)) \geq 2$.

Owing to the above discussion, there is an edge $(u^i, v^i) \in F^i$ such that (1) u^i is incident with a fault-free j-dimension edge; (2) v^i is incident with a fault-free j-dimension edge; (3) $\delta(BH^{j,i}_{n-1} - F^i + (u^i, v^i)) \geq 2$, where $|F^i| = 4n-8$ for $i \in \{0, 1, 2, 3\}$.

Subcase 1.2.1: $|F^0| = 4n-8$.

By induction, there is a Hamiltonian cycle C_0 that contains (u, v) in $BH^{j,0}_{n-1} - F^0 + (u^0, v^0)$. Obviously, $|F \cap E(C_0)| \leq 1$. If $F \cap E(C_0) = 1$, let $(a^0, b^0) = (u^0, v^0)$. If $|F \cap E(C_0)| = 0$, let (a^0, b^0) be an edge such that a^0 (resp., b^0) is incident with a fault-free j-dimension edge. We can represent the cycle C_0 as $\langle u, H_0, a^0, b^0, H'_0, u\rangle$ for $2 \cdot 2^{2n-3} > 4n-5$, (a^0, b^0) exists. Let $(a^0, b^1), (a^1, b^2), (a^2, b^3), (a^3, b^0)$ be fault-free j-dimension edges. By Lemma 2.4, there is a fault-free Hamiltonian path H_i that joins (a^i, b^i) in $BH^{j,i}_{n-1}$ for $i = 1, 2, 3$. The cycle $C = \langle u, H_0, a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H'_0, u\rangle$ (see figure 6) is the desired cycle.



Figure 6: Illustration for subcase 1.2.1 of theorem 3.5

Subcase 1.2.2: $|F^1| = 4n-8$ (or $|F^3| = 4n-8$).

In this subcase, $|F^0| \leq 1$. By induction, there is a Hamiltonian cycle C_1 in $BH^{j,1}_{n-1} - F^1 + (u^1, v^1)$ that contains (u^1, v^1), where u^1 (resp., v^1) is incident with a fault-free j-dimension edge. We can assume that $(u^1, v^1) = (a^1, b^1)$ and (b^1, a^0), (a^1, b^2) are fault-free j-dimension edges incident with a^1 or b^1. Let $H_1 = C_1 - (a^1, b^1)$. Since $|F^0| \leq 1$ and $|F \cap \partial D_j| \leq 3 - |F^0|$, we can choose two fault-free edges $(a^0, b^0), (a^0, d^0) \in BH^{j,0}_{n-1} - F^0$ such that b^0 is incident with a fault-free j-dimension edge (b^0, a^3). Particularly, if $a^0 = u$, let $d^0 = v$. Let $(F^0)' = F^0 \cup T$ where $T = \{(a^0, x) \mid x \in N_{BH^{j,0}_{n-1}}(a^0) \setminus \{b^0, d^0\}\}$. Then, $|(F^0)'| \leq 2n-3 \leq 4n-9$ and $\delta(BH^{j,0}_{n-1} - (F^0)') \geq 2$. By induction, there is a Hamiltonian cycle C_0 in $BH^{j,0}_{n-1}$ that contains (u, v). Since a^0 is incident with exactly two fault-free edges in $BH^{j,0}_{n-1} - (F^0)'$, we have $(a^0, b^0) \in E(C_0)$. We represent C_0 as $\langle u, H_0, a^0, b^0, H'_0, u\rangle$. Let (a^2, b^3) be the fault-free j-dimension edge. By Lemma 2.4, there is a fault-free Hamiltonian path H_i in $BH^{j,i}_{n-1}$ that joins a^i and b^i for $i = 2, 3$. Hence, the cycle $C = \langle u, H_0, a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H'_0, u\rangle$ (see figure 7) is the desired cycle.



Figure 7: Illustration for subcase 1.2.2 of theorem 3.5

Subcase 1.2.3: $|F^2| = 4n-8$.

By induction, there is a Hamiltonian in $BH^{j,2}_{n-1} - F^2 + (u^2, v^2)$ that contains (u^2, v^2) and u^2 (resp., v^2) is incident with a fault-free j-dimension edge. We can assume that $(u^2, v^2) = (a^2, b^2)$ and (a^2, b^3), (a^1, b^2) are fault-free j-dimension edges. Let $H_2 = C_2 - (a^2, b^2)$. By induction, there is a fault-free

Hamiltonian cycle C_0 in $BH_{n-1}^{j,0}$ that contains (u,v). Similar to the analysis of subcase 1.1.1.3, we can represent C_0 as $\langle u, H_0, c^{2i-1}, c^{2i}, H_0', u\rangle$ where c^{2i-1} (resp., c^{2i}) is incident with a fault-free j-dimension edge (c^{2i-1}, b^1) (resp., (c^{2i}, b^3)). By Lemma 2.4, there is a fault-free Hamiltonian path H_i of $BH_{n-1}^{j,i}$ that contains (a^i, b^i) for $i = 1, 3$. Hence, the cycle $C = \langle u, H_0, c^{2i-1}, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H^3, a^3, c^{2i}, H_0', u\rangle$ (see figure 8) is the desired cycle.



Figure 8: Illustration for subcase 1.2.3 of theorem 3.5

Subcase 1.3: There is an integer i in $\{0,1,2,3\}$ such that $|F^i| = 4n-7$.

We prove this subcase according to two cases below.

Subcase 1.3.1 $\delta(BH_{n-1}^{j,i} - F^i) \geq 2$ for all $i \in \{0,1,2,3\}$.

Since $4n-7-2(2n-4) = 1$, there are at least three disjoint faulty edges. There are two faulty edges $(a^i, b^i), (c^i, d^i)$ such that a^i, b^i, c^i, d^i are incident with four disjoint fault-free j-dimension edges owing to $|F \cap \partial D_j| = 2$.

Subcase 1.3.1.1: $|F^0| = 4n-7$.

By induction, there is a Hamiltonian cycle C_0 in $BH_{n-1}^{j,0} - F + (a^0, b^0) + (c^0, d^0)$ that contains (u,v). If $|C_0 \cap \{(a^0, b^0), (c^0, d^0)\}| \leq 1$, according to subcase 1.2.1, there is a desired cycle. If $|C_0 \cap \{(a^0, b^0), (c^0, d^0)\}| = 2$, represent C_0 as $\langle u, H_0, a^0, b^0, H_0', c^0, d^0, H_0'', u\rangle$ (or $\langle u, H_0, a^0, b^0, H_0', d^0, c^0, H_0'', u\rangle$). Let $(a^i, b^{i+1}), (c^i, d^{i+1})$ be fault-free j-dimension edges such that $a^i \neq c^i, b^i \neq d^i$ for all $i = 0,1,2,3$. By Lemma 2.6, there are two disjoint paths H_i, H_i' such that (1) H_i joins a^i and b^i, H_i' joins c^i and d^i; (2) $V(H_i \cup H_i') = V(BH_{n-1}^{j,i})$ for $i = 1,2,3$. Hence, the cycle $C = \langle u, H_0, a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0', c^0, d^1, H_1', c^1, d^2, H_2', c^2, d^3, H_3', c^3, d^0, H_0'', u\rangle$(or the cycle $C = \langle u, H_0, a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0', d^0, c^3, H_3', d^3, c^2, H_2', d^2, c^1, H_1', d^1, c^0, H_0'', u\rangle$) (see figure 9) is the desired cycle.

Subcase 1.3.1.2: $|F^1| = 4n-7$ (or $|F^3| = 4n-7$).

By induction, there is a Hamiltonian cycle C_1 in $BH_{n-1}^{j,1} - F + (a^1, b^1) + (c^1, d^1)$ that contains (a^1, b^1). If $(c^1, d^1) \notin C_1$, according to subcase 1.2.2, there is a desired cycle. Thus, assume that $(c^1, d^1) \in C_1$. We can assume that $C_1 = \langle a^1, b^1, H_1, d^1, c^1, H_1', a^1\rangle$ (or $C_1 = \langle a^1, b^1, H_1, c^1, d^1, H_1', a^1\rangle$) and that $(a^0, b^1), (c^0, d^1), (a^1, b^2), (c^1, d^2)$ are fault-free j-dimension edges. Note that $a^0 \neq c^0$. Then, $a^0 \neq u$ or $c^0 \neq u$. Without loss of generality, we assume that $c^0 \neq u$. If there is a vertex x^0 in $BH_{n-1}^{j,0}$ incident with two faulty j-dimension edges, let $(F^0)' = E(BH_n) \cap \{(a^0, x^0), (c^0, x^0)\}$. Otherwise, let $(F^0)' = \{(c^0, \alpha) \mid \alpha \in BH_{n-1}^{j,0}$, where α is incdent with a faulty $j-$dimension edge$\}$. Since $|F \cap \partial D_j| = 2$, $|(F^0)'| \leq 2$. By induction, there is a Hamiltonian cycle C_0 in $BH_{n-1}^{j,0} - (F^0)'$ that contains (u,v). We may represent it as $\langle u, H_0, a^0, b^0, H_0', c^0, d^0, H_0'', u\rangle$. Let $(b^0, a^3), (d^0, c^3), (a^2, b^3), (c^2, d^3)$ be fault-free j-dimension edges where $a^2 \neq c^2, b^3 \neq d^3, a^3 \neq c^3$ (when $C_1 = \langle a^1, b^1, H_1, c^1, d^1, H_1', a^1\rangle$, let $(b^0, a^3), (d^0, c^3), (a^2, d^3), (c^2, b^3)$ be fault-free j-dimension edges). By Lemma 2.6, there are two disjoint paths H_i, H_i' such that (1) H_i joins a^i and b^i, and H_i' joins c^i and d^i; (2) $V(H_i \cup H_i') = V(BH_{n-1}^{j,i})$ for $i = 2,3$. Hence, The cycle $C = \langle u, H_0, a^0, b^1, H_1, d^1, c^0, H_0', b^0, a^3, H_3, b^3, a^2, H_2, b^2, a^1, H_1', c^1, d^2, H_2', c^2,$



Figure 9: Illustration for subcase 1.3.1.1 of theorem 3.5

$d^3, H'_3, c^3, d^0, H''_0, u\rangle$ (or $C = \langle u, H_0, a^0, b^1, H_1, c^1, d^2, H'_2, c^2, b^3, H_3, a^3, b^0, H'_0,\ c^0, d^1, H'_1, a^1, b^2, H_2, a^2,$ $d^3, H'_3, c^3, d^0, H''_0, u\rangle$) (see figure 10) is the desired cycle.



Figure 10: Illustration for subcase 1.3.1.2 of theorem 3.5

Subcase 1.3.1.3: $|F^2| \leq 4n-7$.

By induction, there is a Hamiltonian cycle C_2 in $BH^{j,2}_{n-1} - F + (a^2, b^2) + (c^2, d^2)$ that contains (a^2, b^2). If $(c^2, d^2) \notin E(C_2)$, according to subcase 1.2.3, there is a desired cycle. Now, let $(c^2, d^2) \in E(C_2)$. We can assume that $C_2 = \langle a^2, b^2, H_2, d^2, c^2, H'_2, a^2\rangle$ (or $C_2 = \langle a^2, b^2, H_2, c^2, d^2, H'_2, a^2\rangle$). Let $(a^1, b^2), (c^1, d^2), (a^2, d^3), (c^2, b^3)$ be fault-free j-dimension edges, where $a^1 \neq c^1, b^3 \neq d^3$. By induction, there is a Hamiltonian cycle C_0 that contains (u, v). Since $|F \cap \partial D_j| = 2$, there are two edges $(a^0, b^0), (c^0, d^0) \in E(C_0)$ such that $(a^0, b^1), (c^0, d^1), (b^0, a^3), (d^0, c^3)$ are fault-free j-dimension edges, where $b^1 \neq d^1, a^3 \neq c^3$. We can represent C_0 as $\langle u, H_0, a^0, b^0, H'_0, c^0, d^0, H''_0, u\rangle$. By Lemma 2.6, there are two disjoint paths H_i, H'_i such that (1) H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)



Figure 11: Illustration for subcase 1.3.1.3 of theorem 3.5

$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 1, 3$. Thus, the cycle $\langle u, H_0, a^0, b^1, H_1, a^1, b^2, H_2, d^2, c^1, H'_1, d^1, c^0, H'_0, b^0, a^3, H_3, b^3, c^2, H'_2, a^2, d^3, H'_3, c^3, d^0, H''_0, u\rangle$(or $C = \langle u, H_0, a^0, b^1, H_1, a^1, b^2, H_2, c^2, b^3, H_3, a^3, b^0, H'_0, c^0, d^1, H'_1, c^1, d^2, H'_2, a^2, d^3, H'_3, c^3, d^0, H''_0, u\rangle$) (see figure 11) is the desired cycle.

Subcase 1.3.2: $\delta(BH^{j,i}_{n-1} - F^i) = 1$ for some $i \in \{0, 1, 2, 3\}$.

By Lemma 3.2, there is no more than one 1-rescuable vertex in $BH^{j,i}_{n-1}$, say, w. Suppose that (w, β) is the fault-free edge incident with w. Without loss of generality, we can assume that w is a white vertex. Since $2n - 3 \geq 3$, w is incident with at least three faulty edges in $BH^{j,i}_{n-1}$. Then, there are two faulty edges $(w, b^i), (w, d^i)$ such that b^i, d^i are incident with two disjoint fault-free j-dimension edges owing to $|F \cap \partial D_j| = 2$.

Subcase 1.3.2.1: $|F^0| = 4n - 7$.

By induction, there is a Hamiltonian cycle C_0 in $BH^{j,0}_{n-1} - F + (w, b^0) + (w, d^0)$ that contains (u, v). If $|C_0 \cap \{(w, b^0), (w, d^0)\}| \leq 1$, according to subcase 1.2.1, there is a desired cycle. Now, we can assume that $|C_0 \cap \{(w, b^0), (w, d^0)\}| = 2$.

(a) Suppose that w is incident with two fault-free j-dimension edges. In this instance, we can repre-



Figure 12: Illustration for subcase 1.3.2.1 (a) of theorem 3.5

sent C_0 as $\langle u, H_0, b^0, w, d^0, H'_0, v, u\rangle$. Let (w, b^1), (w, d^1), (a^1, b^2), (c^1, d^2), (a^2, b^3), (c^2, d^3), (a^3, b^0), (c^3, d^0)

be fault-free j-dimension edges. By Lemma 2.6, there are two paths H_i and H'_i such that (1)H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 1, 2, 3$. Thus, the cycle $\langle u, H_0, b^0, a^3, H_3, b^3, a^2, H_2, b^2, a^1, H_1, b^1, w, d^1, H'_1, c^1, d^2, H'_2, c^2, d^3, H'_3, c^3, d^0, H'_0, u\rangle$ (see figure 12) is the desired cycle.

(b) Suppose that w is incident with exactly one fault-free j-dimension edge (w, b^1).

In this instance, we can represent C_0 as $\langle u, H_0, \beta, \gamma, H'_0, b^0, w, d^0, H''_0, u\rangle$. Suppose that γ is incident with a fault-free j-dimension edge (γ, d^1) where $b^1 \neq d^1$. Let (b^0, a^3), (d^0, c^3) and $(a^i, b^{i+1}), (c^i, d^{i+1})$ be fault-free j-dimension edges for $i = 1, 2$. By Lemma 2.6, there are two paths H_i and H'_i such that (1)H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 1, 2, 3$. Thus, the cycle $C = \langle u, H_0, \beta, w, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H'_0, \gamma, d^1, H'_1, c^1, d^2, H'_2, c^2, d^3, H'_3, c^3, d^0, H''_0, u\rangle$ (see figure 13) is the desired cycle.

Suppose that the unique fault-free j-dimension edge incident with γ is (γ, b^1). Let (b^0, a^3), (d^0, c^3) and $(a^i, b^{i+1}), (c^i, d^{i+1})$ be fault-free j-dimension edges for $i = 1, 2$. By Lemma 2.6, there are two paths H_i and H'_i in $BH^{j,i}_{n-1}$ such that (1)H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 2, 3$. By Lemma 2.7, there is a Hamiltonian path in $BH^{j,1}_{n-1} - \{\alpha\}$ that joins a^1 and c^1. Then, the cycle $C = \langle u, H_0, \beta, w, b^1, \gamma, H'_0, b^0, a^3, H_3, b^3, a^2, H_2, b^2, a^1, H_1, c^1, d^2, H'_2, c^2, d^3, H'_3, c^3, d^0, H''_0, u\rangle$ (see figure 13) is the desired cycle.



Figure 13: Illustration for subcase 1.3.2.1 (b) of theorem 3.5

Subcase 1.3.2.2: $|F^i| = 4n - 7$ for $i = 1, 2, 3$.

By induction, there is a Hamiltonian cycle C_i in $BH^{j,i}_{n-1} - F + (w, b^i) + (w, d^i)$ that contains (w, β). Then, $|C^i \cap \{(w, b^i), (w, d^i)\}| \leq 1$. According to subcase 1.2.2 or 1.2.3, there is a desired cycle.

Case 2: $e \in \partial D_j$.

If $|F \cap \partial D_j| = 2$ or $\delta(BH_n - F - \partial D_j) = 1$, by Lemma 3.2, there is an integer $j' \in \{0, 1, \cdots, n-1\} \setminus \{j\}$ such that $|F \cap \partial D_{j'}| \geq 2$ and there is no isolated vertex and no more than one 1-rescuable vertex in $BH_n - \partial D_{j'}$. Note that $e \in BH_n - \partial D_{j'}$. According to case 1, there is a desired cycle. Thus, we only need to show that there is a Hamiltonian cycle in $BH_n - F$ that contains e when $\delta(BH_n - F - \partial D_j) \geq 2$ and $|F \cap \partial D_j| \geq 3$. Without loss of generality, we can assume that $u = a^0, v = b^0$.

Subcase 2.1: $|F^i| \leq 4n - 9$ for all $i = 0, 1, 2, 3$.

Since $2(2n - 3) + 2 > 4n - 5$, we have $|F \cap BH^{j,0}_{n-1}| \leq 2n - 4$ or $|F \cap BH^{j,1}_{n-1}| \leq 2n - 4$. Without loss of generality, we can assume that $|F \cap BH^{j,0}_{n-1}| \leq 2n - 4$. By Lemma 3.3, there is a fault-free j-dimension edge (a^1, b^2) (resp., $((a^2, b^3), (a^3, b^0))$) such that $(a^1, v) \in E(BH_n)$ (resp., $(a^2, b^2), (a^3, b^3) \in E(BH_n)$). By Lemma 2.4, there is a Hamiltonian path H_0 that joins u and b^0. By induction, there is a Hamiltonian cycle C_i in $(BH^{j,i}_{n-1} - F) \cup \{(a^i, b^i)\}$ that contains (a^i, b^i) for $i = 1, 2, 3$. Let

$H_i = C_i - (a^i, b^i)$. The cycle $C = \langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 14) is the desired cycle.



Figure 14: Illustration for subcase 2.1 of theorem 3.5

Subcase 2.2: There is an $i \in \{0, 1, 2, 3\}$ such that $|F^i| = 4n - 8$.
Subcase 2.2.1: $|F^0| = 4n - 8$ (or $|F^1| = 4n - 8$).
Subcase 2.2.1.1: Suppose that u is incident with at least two faulty edges in $BH^{j,0}_{n-1}$.

Since $|F \cap \partial D_j| = 3$ and each vertex is incident with exactly two j-dimension edges, there is a faulty edge (u, b^0) such that b^0 is incident with a fault-free j-dimension edge, say, (a^3, b^0). By induction, there is a Hamiltonian cycle C_0 in $BH^{j,0}_{n-1}$ that contains (u, b^0). Let $H_0 = C_0 - (u, b^0)$ and $(a^1, b^2), (a^2, b^3)$ be fault-free j-dimension edges. By Lemma 2.6, there is a path H_i in $BH^{j,i}_{n-1}$ that joins a^i and b^i. Therefore, the cycle $\langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 15) is the desired cycle.



Figure 15: Illustration for subcase 2.2.1.1 of theorem 3.5

Subcase 2.2.1.2: Suppose that u is incident with no more than one faulty edge in $BH^{j,0}_{n-1}$.

In this subcase, there are two disjoint faulty edges in $BH^{j,0}_{n-1} - \{u\}$, say, $(c^0, d^0), (x^0, y^0)$, that are the result of $4n - 8 - (2n - 4) - 1 = 2n - 5 \geq 1$. Suppose that x^0, a^0 is a white vertex and $M = \{z \mid (x^0, z) \in \partial D_j \setminus F$ or $(c^0, z) \in \partial D_j \setminus F\}$.
Subcase 2.2.1.2.1: $M \neq \{v\}$.

Without loss of generality, we can assume that (c^0, d^1) is a fault-free j-dimension edge where $d^1 \neq v$.

(a) Suppose that d^0 is incident with a fault-free j-dimension edge (d^0, c^3).

Note that u is incident with no more than one faulty edge in $BH^{j,0}_{n-1}$. There is a vertex $b^0 \in N_{BH^{j,0}_{n-1}}(u)$ such that b^0 is incident with a fault-free j-dimension edge (b^0, a^3) where $a^3 \neq c^3$ owing to $2(2n-3)-3 \geq 3$. By induction, there is a Hamiltonian cycle C_0 in $(BH^{j,0}_{n-1} - F + (c^0, d^0)) \cup \{(u, b^0)\}$ that contains (u, b^0). If $(c^0, d^0) \notin E(C_0)$, similar to subcase 2.2.1.1, there is a desired cycle. Thus, we can assume that $(c^0, d^0) \in E(C_0)$. We can represent C_0 as $\langle u, b^0, H_0, c^0, d^0, H'_0, u\rangle$. Let $(a^i, b^{i+1}), (c^i, d^{i+1})$ be fault-free j-dimension edges for $i = 1, 2$ where $a^i \neq c^i, b^i \neq d^i$. By Lemma 2.6, there are two vertex-disjoint paths H_i and H'_i such that (1)H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 1, 2, 3$. Hence, the cycle $C = \langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, c^0, d^1, H'_1, c^1, d^2, H'_2, c^2, d^3, H'_3, c^3, d^0, H'_0, u\rangle$ (see figure 16 (a)) is the desired cycle.

(b) Suppose that d^0 is incident with two faulty j-dimension edges.

By induction, there is a Hamiltonian cycle in $BH^{j,0}_{n-1} - F + (c^0, d^0)$ that contains (c^0, d^0). Since $\delta(BH_n - F - \partial D_j) \geq 2$, d^0 is incident with a fault-free edge (d^0, α) in $BH^{j,0}_{n-1}$, which does not belong to $E(C_0)$. Without loss of generality, we can represent C_0 as $\langle u, H_0, d^0, c^0, H'_0, \gamma, \alpha, H''_0, \beta, u\rangle$. Note that d^0 is incident with two faulty j-dimension edges, and β, γ are incident with two disjoint fault-free j-dimension edges $(\beta, c^3), (\gamma, a^3)$. Let $(a^i, b^{i+1}), (c^i, d^{i+1})$ be fault-free j-dimension edges where $a^i \neq c^i, b^i \neq d^i$ for $i = 1, 2$. By Lemma 2.6, there are two Hamiltonian paths H_i, H'_i such that (1)H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 1, 2, 3$. Thus, the cycle $\langle u, H_0, d^0, \alpha, H''_0, \beta, c^3, H'_3, d^3, c^2, H'_2, d^2, c^1, H'_1, d^1, c^0, H'_0, \gamma, a^3, H_3, b^3, a^2, H_2, b^2, a^1, H_1, v, u\rangle$ (see figure 16 (b)) is the desired cycle.



Figure 16: Illustration for subcase 2.2.1.2.1 (a) and (b) of theorem 3.5

Subcase 2.2.1.2.2: $M = \{v\}$.

In this subcase, $|E(BH^{j,0}_{n-1}, BH^{j,1}_{n-1}) \cap F| = 3$ and $|E(BH^{j,0}_{n-1}, BH^{j,3}_{n-1}) \cap F| = 0$. By induction, there is a Hamiltonian cycle C_0 in $BH^{j,0}_{n-1} - F + (c^0, d^0)$ that contains (c^0, d^0). We can represent C_0 as $\langle u, b^0, H_0, c^0, d^0, H'_0, u\rangle$. Since b^0 is incident with two fault-free j-dimension edges, we can choose a fault-free j-dimension edge (b^0, a^3) where $a^3 \neq c^3$. Let $(a^i, b^{i+1}), (c^i, d^{i+1})$ be fault-free j-dimension edges for $i = 1, 2$ where $a^i \neq c^i, \neq b^i \neq d^i$. By Lemma 2.6, there are two vertex-disjoint paths H_i and H'_i such that (1)H_i joins a^i and b^i, and H'_i joins c^i and d^i; (2)$V(H_i \cup H'_i) = V(BH^{j,i}_{n-1})$ for $i = 2, 3$. By Lemma 2.7, there is a Hamiltonian path H_1 in $BH^{j,1}_{n-1} - \{v\}$ that joins a^1 and c^1. Hence, the cycle $C = \langle u, v, c^0, H_0, b^0, a^3, H_3, b^3, a^2, H_2, b^2, a^1, H_1, c^1, d^2, H'_2, c^2, d^3, H'_3, c^3, d^0, H'_0, u\rangle$ (see figure 17) is the desired cycle.

Subcase 2.2.2: $|F^2| = 4n-8$ (or $|F^3| = 4n-8$).

There are two disjoint faulty edges in $BH^{j,2}_{n-1}$ for $4n-8-(2n-4) = 2n-4 \geq 2$. Since $|F \cap \partial D_j| = 3$, there is a faulty edge $(a^2, b^2) \in BH^{j,2}_{n-1}$ such that a^2 (resp., b^2) is incident with a fault-free j-dimension edge (a^2, b^3) (resp., (a^1, b^2)). By induction there is a Hamiltonian cycle C_2 in $BH^{j,2}_{n-1} - F + (a^2, b^2)$ that contains (a^2, b^2). Let $H_2 = C_2 - (a^2, b^2)$ and (a^3, b^0) be a fault-free j-dimension edge. By Lemma



Figure 17: Illustration for subcase 2.2.1.2.2 of theorem 3.5

2.4, there is a Hamiltonian path H_i in $BH^{j,i}_{n-1}$ that joins a^i and b^i for $i = 0, 1, 3$. Thus, the cycle $C = \langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 18) is the desired cycle.



Figure 18: Illustration for subcase 2.2.2 of theorem 3.5

Appendix A. Proof of Lemma 3.4

Lemma 3.4 Let $F \subseteq E(BH_2)$ with $|F| \leq 3$ and $\delta(BH_2 - F) \geq 2$. Then, each edge in $BH_2 - F$ lies on a fault-free Hamiltonian cycle.

Suppose that $|F| = 3$. By Lemma 2.3, without loss of generality, we can assume that $|F \cap \partial D_1| \geq |F \cap \partial D_0|$. Let $e = (u, v)$. In the following, we can assume that a^i, c^i are white vertices and b^i, d^i are black vertices in $BH^{1,i}_1$ for $i = 0, 1, 2, 3$.

Case 1: $|F \cap \partial D_1| = 3$, $|F \cap \partial D_0| = 0$.

Subcase 1.1: $e \in \partial D_0$.

Without loss of generality, we can assume that $e \in BH^{1,0}_1$. Since $|F \cap \partial D_1| = 3$, there is an edge (x, y) in $\{(v, a^0), (u, b^0)\}$ such that x (resp., y) is incident with a fault-free 1-dimension edge. Without loss of generality, we can assume that $(x, y) = (u, b^0)$. Let $H_0 = BH^{1,0}_1 - (u, b^0)$ and $(u, b^1), (b^0, a^3), (a^1, b^2), (a^2, b^3)$ be fault-free 1-dimension edges. Note that $|F \cap \partial D_0| = 0$. There is a Hamiltonian path H_i in $BH^{1,i}_1$ that joins a^i and b^i for $i = 1, 2, 3$. Thus, the cycle $C = \langle u, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b_0, H_0, u\rangle$ (see figure 19)(a) is the desired cycle.

Subcase 1.2: $e \in \partial D_1$.

Without loss of generality, we can assume that (u, v) is an edge between $BH_1^{1,0}$ and $BH_1^{1,1}$ and $u = a^0, v = b^1$. Let $(a^1, b^2), (a^2, b^3), (a^3, b^0)$ be fault-free 1-dimension edges. Since $|F \cap \partial D_0| = 0$, there is a Hamiltonian path H_i of $BH_1^{1,i}$ that joins a^i and b^i for $i = 0, 1, 2, 3$. Thus, the cycle $\langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 19 (b)) is the desired cycle.

Case 2: $|F \cap \partial D_1| = 2$, $|F \cap \partial D_0| = 1$.

Subcase 2.1: $e \in \partial D_0$.

Without loss of generality, we can assume that $e \in BH_1^{1,0}$.

Subcase 2.1.1: $|F^0| = 1$.

Suppose that (a^0, b^0) is a faulty edge in $BH_1^{1,0}$. Let $H_0 = BH_1^{1,0} - (a^0, b^0)$. Then, H_0 is a fault-free Hamiltonian path in $BH_1^{1,0}$ that contains (u, v). Since $\delta(BH_2 - F) \geq 2$, a^0 (resp., b^0) is incident with a fault-free 1-dimension edge (a^0, b^1) (resp., (b^0, a^3)). Let $(a^1, b^2), (a^2, b^3)$ be fault-free 1-dimension edges. Since $|F^1| = |F^2| = |F^3| = 0$, there is a Hamiltonian path H_i in $BH_1^{1,i}$ that joins a^i and b^i for $i = 1, 2, 3$. Thus, the cycle $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0\rangle$ (see figure 19 (c))is the desired cycle.

Subcase 2.1.2: $|F^1| = 1$(or $|F^3| = 1$).

Let $V(BH_1^{1,0}) = \{u, v, a^0, b^0\}$ and (a^1, b^1) be the faulty edge in $BH_1^{1,1}$. Suppose that $H_1 = BH_1^{1,1} - (a^1, b^1)$. Then, H_1 is a Hamiltonian path in $BH_1^{1,1}$ that joins a^1 and b^1. Since $\delta(BH_2 - F) \geq 2$, a^1 (resp., b^1) is incident with a fault-free 1-dimension edge.

(a) $(b^1, a^0) \in F$. Since $|F \cap \partial D_1| = 2$ and $(b^1, a^0) \in F$, b^0 is incident with at least one fault-free 1-dimension edge, say, (b^0, a^3). Let $H_0 = \langle b^0, a^0, v, u\rangle$ and $(a^1, b^2), (a^2, b^3)$ be fault-free 1-dimension edges. Since $|F^2| = |F^3| = 0$, there is a Hamiltonian path H_i in $BH_1^{1,i}$ that joins a^i and b^i for $i = 2, 3$. Hence the cycle $C = \langle u, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 19 (d)) is the desired cycle.

(b) $(b^1, a^0) \notin F$. Since $|F \cap D_1| = 2$, v or b^0 is incident with a fault-free 1-dimension edge. Without loss of generality, we can assume that b^0 is incident with a fault-free 1-dimension edge (b^0, a^3). Let $H_0 = \langle b^0, u, v, a^0\rangle$ and $(a^1, b^2), (a^2, b^3)$ be fault-free 1-dimension edges. Since $|F^2| = |F^3| = 0$, there is a Hamiltonian path H_i in $BH_1^{1,i}$ that joins a^i and b^i for $i = 2, 3$. Thus, the cycle $C = \langle b^0, H_0, a^0, b_1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0\rangle$ (see figure 19 (e)) is the desired cycle.

Subcase 2.1.3: $|F^2| = 1$.

We can represent $BH_1^{1,0}$ as $\langle u, v, a^0, b^0, u\rangle$. Since $|F \cap \partial D_1| = 2$, we can see that there is an edge (x, y) in $\{(v, a^0), (u, b^0)\}$ such that x (resp., y) is incident with a fault-free j-dimension edge. Without loss of generality, we can assume that $(x, y) = (u, b^0)$ and $(u, b^1), (b^0, a^3)$ are fault-free 1-dimension edges. Let $H_1 = \langle b^0, a^0, v, u\rangle$ and (a^2, b^2) be the faulty edge in $BH_1^{1,2}$. Since $\delta(BH_2 - F) \geq 2$, a^2 (resp., b^2) is incident with a fault-free 1-dimension edge (a^2, b^3) (resp., (a^1, b^2)). Note that $|F^1| = |F^3| = 0$. There is a Hamiltonian path H_i in $BH_1^{1,i}$ that joins a^i and b^i for $i = 1, 3$. Thus, the cycle $C = \langle u, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 19 (f)) is the desired cycle.

Subcase 2.2: $e \in \partial D_1$.

Without loss of generality, we can assume that $e = (u, v) = (a^0, b^1)$.

Subcase 2.2.1: $|F^0| = 1$(or $|F^1| = 1$).

Suppose that $V(BH_1^{1,i}) = \{a^i, b^i, c^i, d^i\}$.

Subcase 2.2.1.1: (a^0, x) is a faulty edge, where $x \in \{b^0, d^0\}$.

Without loss of generality, we can assume that $x = b^0$. Since $\delta(BH_2 - F) \geq 2$, b^0 is incident with a fault-free 1-dimension edge, say, (b^0, a^3). Let $H_0 = BH_1^{1,0} - (a^0, b^0)$ and $(a^1, b^2), (a^2, b^3)$ be fault-free 1-dimension edges. Note that $|F^1| = |F^2| = |F^3| = 0$. There is a Hamiltonian path H_i of $BH_1^{1,i}$ that joins a^i and b^i for $i = 1, 2, 3$. Thus, the cycle $C = \langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 19 (g)) is the desired cycle.

Subcase 2.1.1.2: (c^0, x) is a faulty edge where $x \in \{b^0, d^0\}$.

Without loss of generality, we can assume that $x = b_0$.

(a) Suppose that each vertex in BH_2 is incident with at least one fault-free 1-dimension edge. Then, there are two disjoint fault-free 1-dimension edges between $BH_1^{1,i}$ and $BH_1^{1,i+1}$ for all $i = 1, 2, 3$, namely $(a^i, b^{i+1}), (c^i, d^{i+1})$ for $i = 1, 2, 3$. Since $\delta(BH_2 - F) \geq 2$ and $(b^0, c^0) \in F$, (c^0, b^1) or (c^0, d^1) is a fault-free 1-dimension edge.



Figure 19: Illustration for Lemma 3.4

If $(c^0, d^1) \notin F$, the cycle $\langle u, v, c^0, d^0, c^3, d^3, c^2, d^2, c^1, d^1, a^1, b^2, a^2, b^3, a^3, b^0, u\rangle$ (see figure 19 (h)) is the desired cycle.

If $(c^0, d^1) \notin F$, the cycle $\langle u, v, c^1, d^2, c^2, d^3, c^3, d^0, c^0, d^1, a^1, b^2, a^2, b^3, a^3, b^0, u\rangle$ (see figure 19 (i)) is the desired cycle.

(b) Suppose that there is a vertex y such that y is incident with two faulty 1-dimension edges. Without loss of generality, we can let $y = a^1$. Then, the cycle $C = \langle u, v, a^1, d^1, c^0, d^0, c^3, d^3, c^2, d^2, c^1, b^2, a^2, b^3, a^3, b^0, u\rangle$ (see figure 19 (j)) is the desired cycle.

Subcase 2.2.2: $|F^2| = 1$(or $|F^3| = 1$).

Let (a^2, b^2) be the faulty edge in $BH_1^{1,2}$. Since $\delta(BH_2 - F) \geq 2$, a^2 (resp., b^2) is incident with a fault-free 1-dimension edge (a^2, b^3) (resp., (a^1, b^2)). Since $|F \cap \partial D_1| = 2$, there is a fault-free 1-dimension edge incident with b^0 or d^0. Without loss of generality, we can assume that (b^0, a^3) is a fault-free 1-dimension edge. Let $H_i = BH_1^{1,i} - (a^i, b^i)$ for $i = 0, 1, 2, 3$. Then, the cycle $C = \langle u, v, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u\rangle$ (see figure 19 (k)) is the desired cycle.

4 Conclusion

In this paper, we consider the problem proposed by Zhou [25] and show that each fault-free edge lies on a fault-free Hamiltonian cycle after no more than $4n-5$ faulty edges occur when each vertex is incident with at least two fault-free edges. Our result is optimal because there is a counterexample when $|F| = 4n - 4$ as follows. Let $u = (0, 0, \cdots, 0), v = (2, 0, \cdots, 0), x = (1, 0, \cdots, 0), y = (3, 0, \cdots, 0)$ and $F = \{(u, a) \mid a \in N_{BH_n}(u) \setminus \{x, y\}\} \bigcup \{(v, b) \mid b \in N_{BH_n}(u) \setminus \{x, y\}\}$. Then, $|F| = 4n-4$. Suppose that there is a Hamiltonian cycle C. Note that $u, v \in V(C)$. Then, C contains $(u, x), (u, y), (v, x), (v, y)$. Note that $\langle u, x, v, y, u\rangle$ is a cycle of length 4, which is a contradiction. Thus, there is no Hamiltonian cycle in $BH_n - F$.

Acknowledgement

This research is supported by the National Natural Science Foundation of China (11571044, 61373021), the Fundamental Research Funds for the Central University of China.

References

[1] J.A. Bondy, U.S.R. Murty, Graph theory with applitions. *North Holland, New York*, 1980.

[2] J.C. Chen, C.H. Tsai, Conditional edge-fault-tolerant Hamiltonicity of dual-cubes, *Inform. Sci.* 181 (2011) 620-627.

[3] D.Q. Cheng, R.X. Hao, Y.Q. Feng, Vertex-fault-tolerant cycles embedding in balanced hypercubes, *Inform. Sci.* 288 (2014) 449-461.

[4] D.Q. Cheng, R.X. Hao, Y.Q. Feng, Two node-disjoint paths in balanced hypercubes, *Appl. Math. Comput.* 242(2014) 127-142.

[5] D.Q. Cheng, R.X. Hao, Various cycles embedding in faulty balanced hypercubes, *Inform. Sci.* 297 (2015) 140-153.

[6] R.X. Hao, R. Zhang, Y.Q. Feng, J.X. Zhou, Hamiltonian cycle embedding for fault tolerant in balanced hypercubes, *Appl. Math. Comput.* 244 (2014) 447-456.

[7] S.Y. Hsieh, Y.R. Cian, Conditional edge-fault Hamiltonicity of argmented cubes, *Inform. Sci.* 180 (2010) 2596-2617.

[8] K. Huang, J. Wu, Area efficient layout of balanced hypercubes, *Int. J. High Speed Electron. Syst.* 6(4) (1995) 631-646.

[9] , H.S. Hung, J.S. Fu, G.H. Chen, Fault-free Hamiltonian cycles in crossed cubes with conditional link faults, *Infom. Sci.* 177 (2007) 5664-5674.

[10] J.J. Liu, Y.L. Wang, Hamiltonian cycles in hypercubes with faulty edges, *Inform. Sci.* 256 (2014) 225-233.

[11] H. Lü, X. Li, H. Zhang, Matching preclusion for balanced hypercubes, *Theor. Comput. Sci.* 465 (2012) 10-20.

[12] H.Z. Lü, H.P. Zhang, Hyper-Hamiltonian laceability of balanced hypercubes, *J. Supercomput.* 68 (2014) 302-314.

[13] M.J. Ma, G.Z. Liu, J.M. Xu, Fault-tolerant embedding of paths in crossed cubes, *Theor. Comput. Sci.* 407 (2008) 110-116.

[14] S.N. Sun, M. Xu, K.S. Wang, Edge-fault-tolerant pancyclicity of arrangement graphs, *Inform. Sci.* 285 (2014) 50-62.

[15] H.L. Wang, J.W. Wang, J.M. Xu, Edge-fault-tolerant bipanconnectivity of hypercubes, *Inform. Sci.* 179 (2009) 404-409.

[16] W.Q. Wang, X.B. Chen, A fault-free Hamiltonian cycle passing through prescribed edges in a hypercube with faulty edges, *Inform. Process Lett.* 107 (208) 205-210.

[17] J. Wu, K. Huang, The balanced hypercubes: A cube-based system for fault-tolerant applitions, *IEEE Trans. Comput.* 46(4) (1997) 484-490.

[18] J.M. Xu, M.J. Ma, Z.Z. Du, Edge-fault-tolerant properties of hypercubes and folded hypercubes. *Australasian J. Combinatorics* 35 (2006) 7-16.

[19] J.M. Xu, M.J. Ma, A Survey on Cycle and Path Embedding in Some Networks. *Frontiers of Mathematics in China*, 4 (2) (2009), 217-252.

[20] M. Xu, X.D. Hu, J.M. Xu, Edge-pancyclicity and Hamiltonian laceability of the balanced hypercubes, *Appl. Math. Comput.* 189 (2007) 1393-1401.

[21] D.W. Yang, Y.Q. Feng, J.H. Kwak, J.X. Zhou, Fault-tolerant edge-bipancyclicity of faulty hypercubes under the conditional-fault model, *Inform. Sci.* 329 (2016) 317-328.

[22] M.C. Yang, Bipanconnectivity of balanced hypercubes, *Comput. Math. Appl.* 60 (2010) 1859-1867.

[23] M.C. Yang, Super connectivity of balanced hypercubes, *Appl. Math. Comput.* 219 (2012) 970-975.

[24] M.C. Yang, T.K. Li, J.J.M Tan, L.H. Hsu, On embedding cycles into faulty twisted cubes, *Infom. Sci.* 176 (2006) 676-690.

[25] Q.G. Zhou, C. Chen, H.Z. Lü, Fault-tolerant Hamiltonian laceability of balanced hypercubes, *Inform. Sci.* 300 (2015) 20-27.

[26] J.X. Zhou, Z.L. Wu, S.C Yang, K.W. Yuan, Symmetric property and reliability of balanced hypercube, *IEEE Trans. Comput.* 64(3) (2015) 876-881.